

ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
әл-Фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Ұлттық академиясының
им. әл-Фараби

N E W S

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-Farabi
Kazakh National University

SERIES PHYSICO-MATHEMATICAL

2 (336)

MARCH – APRIL 2021

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series physico-mathematical journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of chemistry and technologies in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of chemical sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Физикалық-математикалық сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Химия және технология сериясы Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді химиялық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия физико-математическая» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по химическим наукам для нашего сообщества.

Б а с р е д а к т о р
ф.-м.ғ.д., проф., ҚР ҰҒА академигі
Ғ.М. Мұтанов

Р е д а к ц и я а л қ а с ы:

Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Байгунчечков Ж.Ж. проф., академик (Қазақстан)
Quevedo Hernando проф. (Мексика)
Жүсіпов М.А. проф. (Қазақстан)
Ковалев А.М. проф., академик (Украина)
Калимолдаев М.Н. проф., академик (Қазақстан)
Михалевич А.А. проф., академик (Белорусь)
Мырзакулов Р. проф., академик (Қазақстан)
Рамазанов Т.С. проф., академик (Қазақстан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)
Уалиев З.Г. проф., чл.-корр. (Қазақстан)
Харин С.Н. проф., академик (Қазақстан)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).
Қазақстан Республикасының Ақпарат және коммуникациялар министрлігінің Ақпарат комитетінде
14.02.2018 ж. берілген № **16906-Ж** мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Тақырыптық бағыты: *физика-математика ғылымдары және ақпараттық
технологиялар саласындағы басым ғылыми зерттеулерді
жариялау.*

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекен-жайы: 050010, Алматы қ., Шевченко көш., 28; 219 бөл.;
тел.: 272-13-19; 272-13-18

<http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2021

Типографияның мекен-жайы: «Аруна» ЖК, Алматы қ., Мұратбаева көш., 75.

Главный редактор
д.ф.-м.н., проф. академик НАН РК
Г.М. Мутанов

Редакционная коллегия:

Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Байгунчеков Ж.Ж. проф., академик (Казахстан)
Quevedo Hernando проф. (Мексика)
Жусупов М.А. проф. (Казахстан)
Ковалев А.М. проф., академик (Украина)
Калимолдаев М.Н. проф., академик (Казахстан)
Михалевич А.А. проф., академик (Беларусь)
Мырзакулов Р. проф., академик (Казахстан)
Рамазанов Т.С. проф., академик (Казахстан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)
Уалиев З.Г. проф., чл.-корр. (Казахстан)
Харин С.Н. проф., академик (Казахстан)

«Известия НАН РК. Серия физика-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации Министерства информации и коммуникаций Республики Казахстан № 16906-Ж, выданное 14.02.2018 г.

Тематическая направленность: *публикация приоритетных научных исследований в области физико-математических наук и информационных технологий.*

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28; ком. 219; тел.: 272-13-19; 272-13-18

<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2021

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

Editor in chief
doctor of physics and mathematics, professor, academician of NAS RK
G.M. Mutanov

Editorial board:

Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Baigunchekov Zh.Zh. prof., akademik (Kazakhstan)
Quevedo Hemando prof. (Mexico)
Zhusupov M.A. prof. (Kazakhstan)
Kovalev A.M. prof., academician (Ukraine)
Kalimoldaev M.N. prof., akademik (Kazakhstan)
Mikhalevich A.A. prof., academician (Belarus)
Myrzakulov R. prof., akademik (Kazakhstan)
Ramazanov T.S. prof., akademik (Kazakhstan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)
Ualiev Z.G. prof., chl.-korr. (Kazakhstan)
Kharin S.N. prof., academician (Kazakhstan)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodical printed publication in the Committee of information of the Ministry of Information and Communications of the Republic of Kazakhstan **No. 16906-Ж**, issued on 14.02.2018.

Thematic scope: *publication of priority research in the field of physical and mathematical sciences and information technology.*

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, Almaty, 050010, tel. 272-13-19; 272-13-18

<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2021

Address of printing house: ST "Aruna", 75, Muratbayev str., Almaty.

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

Volume 2, Number 336 (2021), 39 – 47

<https://doi.org/10.32014/2021.2518-1726.19>

UDC 004.2

M. Sarsembayev¹, B. Urmashev¹, O. Mamyrbayev², M. Turdalyuly³, T. Sarsembayeva¹

¹Al-Farabi Kazakh National University, Almaty, Kazakhstan;

²Institute of Information and Computational Technologies;

³Satbayev University.

E-mail: magjan@kaznu.kz, baidaulet.urmashev@kaznu.kz, morkenj@mail.ru,
m.turdalyuly@satbayev.university, sagdatbek.talshyn1@kaznu.kz

USING THE CUDA TECHNOLOGY TO SPEED UP COMPUTATIONS IN PROBLEMS OF CHEMICAL KINETICS

Abstract. The main idea of the implementation is reducing the time for calculation and thereby implement a multi-user mode for users by placing it on a server with access via a web browser. To model the kinetics of chemical reacting systems were used 4th and 5th grade Runge-Kutta methods and to receive the index of advantages of this elaboration were written programs in C# for sequential computation on a central processor and was used a platform for parallel computation of CUDA on graphic processors. Parallelization of data during calculation on a GPU was performed by the distribution of the reaction to individual strands, when changes of the concentration was calculated over a given time interval of a certain substance. Parallelization is performed over all elementary reactions, with the increasing of the number of reactions in the mechanism, because of this the computation on the GPU has a noticeable gain in time.

Keywords: CUDA, combustion, modeling, parallel computation, performance, computing, thermodynamics, concentration changes, reactions of mechanism.

Nowadays computing technologies significantly permits to increase the speed of mathematical modeling of any processes. However not all program complexes efficiently allows to use parallel processing of the received data. In this case, it concerns to parallel computing with using graphics processors [1-3].

The purpose of the work is to use an NVIDIA graphics card with the calculative abilities of the graphic processors for acceleration of definite chemical reactions processing under different temperature regimes. There was conducted a comparative analysis of programming technologies using graphic processors is carried out. After analyzing of their features and capabilities, the CUDA (Compute Unified Device Architecture) technology, which was developed by NVIDIA, was chosen as the most convenient in terms of programming and presented features. CUDA is a parallel computing architecture from NVIDIA, which significantly allows you to increase computing productivity due to using the GPUs (graphics processors). The applying of this a parallelization technology for solving the equations of chemical kinetics makes it possible to reduce the time and computational costs [4-10].

CUDA computing flows are performed on a separate physical device, the so-called device, which is used as a central processor's (CPU) coprocessor. In its turn, the program in CUDA-C language, the so-called kernel, starts from the central processor for each of the computational flows. The threads' configuration is set on the CPU before the kernel function's calling. For convenience and best performance, computational flows are combined into blocks, which then are combined into a grid. In the course of the implementation of then massively parallel kinetic solver was developed the "Chemical Calculator" software for simulating the kinetics of chemically reacting systems at the zero-dimensional approximation. The work of the program combines the process of computing, the central processing unit and the NVIDIA graphics processor, using CUDA technologies [11-14].

The graphical interface of the program was developed with the use of C # language [2-5]. The program interface was created in the maximally convenient way for users at solving problems of chemically reacting systems. The type of software is shown in figure 1.

Abbreviated files of the Chemkin mechanism are used at the performing of calculations in the developed program. It allows the user to easily transfer their bulk mechanisms into the Chemkin format [2; 4; 6]. The user can download these files using the menu items File → Reactions and File → Thermodynamic Properties. After downloading, the input files appear in the "Input file" program area. Here the separator between the reactions and the thermodynamic properties of substances is a string, which contains the word "THERM". Reading data from the "Input file" area is done line by line and the results of reading are written at the "Reading results" area. The number of reagents and reactions in a given mechanism, the Arrhenius formula coefficients, the direction of the reaction, the derivation of the left and right matrix for the reaction mechanism and the thermodynamic data for each element in the mechanism are determined as a result of reading. The definition of this data is made with only one click of the menu item Reading → Reading the data from a file. After this the input data is entered for conducting calculations, choosing a certain method from the drop-down list. Calculations can be conducted on the central processor as well as on the GPU [6; 10]. At performing of the calculations on NVIDIA with an increasing of the number of computations, time of calculations decreases in comparison with computing time on central processors [11-14].




Figure 1 - The main interface of the program "Chemical calculator"

To simulate the kinetics of chemical reacting systems used of grade Runge-Kutta methods. This methods was written a C # program code for sequential computation on the central processor and CUDA technology was used for parallel computation [14-17].

Table 1 - Comparative analysis of the estimated time in the CPU and GPU for different sizes of the input mechanism

Mechanism: Aramco Mech				
4 th grade of Runge-Kutta method				
Number of chemical elements	Number of chemical reactions	Regime of calculations of CPU, msec	Regime of calculation of GPU, msec	Received accelerationx10
21	51	6901,00	9080,24	0,76
33	118	39081,00	20204,50	1,93
39	191	87735,00	29619,10	2,96
52	245	199169,00	49747,80	4,00
61	301	336749,00	67557,10	4,98
66	376	491811,00	83188,70	5,91
84	448	946439,00	125339,00	7,55
103	565	1795548,00	186175,00	9,64
162	1006	6832052,80	390403,00	17,5

In table 1 is given comparative analysis of CPU and GPU time of calculation. Parallelization of data during the calculation for the GPU is carried out by distributing the reactions on separate strings, at the calculating changes of concentrations in a given time interval of a certain substance [15; 18; 19-28].

Each thread calculates how much the concentration of a given substance in a definite reaction has changed over a certain time interval. Parallelization is performed by reactions with increasing of the number of reactions in the mechanism, that's why computation on the GPU considerably saves time (table 1).

The program implements automated reading of data from input files to the Chemkin software package format, subtracted algorithms for calculating rates of reactions, calculating changes in the concentration of substances over time [25-28]. The main calculations could be performed by 4th grade of Runge-Kutta method [29]. The computation on the central processor is performed by using a sequential algorithm. Calculation on NVIDIA graphics cards, using CUDA parallelization technologies, is performed in parallel way. The use of enterprise software was made by a comparative analysis of the computation time on the central processing unit and on the graphics processor. As a result, it was found that accelerating of the computation process using graphics cards gives a very good time win [27-29].




Figure 2 - Comparative analysis of computational algorithms on CPU and GPU using The fourth-order Runge-Kutta method

User's manual of the program "Chemical calculator". When you start the program, opens the main interface. The program interface consists of several components. In the upper area of the interface is a row of menu components. Loading of input files into the program is conducted through the menu item File. File - Thermodynamic properties (figure 2). These downloaded files are imported into the "Input file" program area. "Thermodynamic properties" - the program for automatic insertion of the word-separator "THERMDAT". During the reading of data from the "Input file" area, the beginning of the thermodynamic data line is determined from this word. If necessary, you can edit the input data of the mechanism directly in this area [25-29].

Reading of data from the input area is performed by clicking the menu item Read → Read data from the file. After clicking this menu item on the created algorithm of data recognition from the input area realizes the automatically reading of the data. At reading the data of reaction, four parameters are determined for calculating their action rates (formulas 1, 2).

The rate of the reverse reaction is determined by the following formula

$$k_{ri} = \frac{k_{fi}}{K_{ci}} \quad (1)$$

$$K_{ci} = K_{pi} \left(\frac{P_{atm}}{RT} \right)^{\sum_{k=1}^K \nu_{ki}} \quad (2)$$

The first parameter is the pre-exponential factor, the second parameter is the temperature exponent, the third parameter is the energy activation and the fourth parameter determines the direction of reaction. If the reaction proceeds only in one direction, then only the rate of reaction is calculated only in this direction, in the other case it depends on the rate of the forward and reverse reactions. After reading these reactions, from reactions they all correspond to the reagents of this mechanism [23-29].

The left and right data matrix is then determined. In these matrices the corresponding coefficients of the elements in the given reaction are located. These data are also used at determining in rates of reaction (formulas 3, 4).

The rate of formation of the k -th substance ($\dot{\omega}_k$) is calculated by the following formula [25-29]:

$$\dot{\omega}_k = \sum_{i=1}^I v_{ki} q_i \quad (k = 1, \dots, K) \quad (3)$$

Here $v_{ki} = v''_{ki} - v'_{ki}$.

Rate of reaction progress' changing of i -reaction determines by following formula:

$$q_i = k_{fi} \prod_{k=1}^K [X_k]^{v''_{ki}} - k_{ri} \prod_{k=1}^K [X_k]^{v'_{ki}} \quad (4)$$

$[X_k]$ – the molar concentration of the k -substance, k_{fi} and k_{ri} the rate of direct and reverse reactions.

The rate of the direct reaction is determined by the Arrhenius form

$$k_{fi} = A_i T^{\beta_i} \exp \left(\frac{-E_i}{R_c T} \right) \quad (5)$$

Where R is the universal gas constant; A_i is a pre-exponential factor that does not depend on temperature, but is determined only by the type of reaction; β_i is the temperature exponent, E_i is the activation energy of the i -th reaction, which can be characterized as a certain threshold energy: roughly speaking, if the energy of the colliding particles is smaller, then the collision will not occur if the energy exceeds E_i . The activation energy does not depend on temperature.




Figure 3 - Reading data from a file

Thermodynamic properties are read for each reagent of this mechanism from the input data of thermodynamic properties. Thermodynamic properties are determined by the heat capacity, enthalpy, entropy, free energies of Gibbs and Helmholtz, for a given substance (formulas 1-5) [28,29].

Basic equations of energy conservation in a medium with constant pressure

$$\frac{dT}{dt} = -\frac{1}{\rho \bar{c}_p} \sum_{k=1}^K H_k \dot{\omega}_k \quad (6)$$

Here, T is the temperature of the medium, ρ is the mass density, $\bar{c}_p$ is the specific heat, H_k is the molar enthalpy of substances and $\dot{\omega}_k$ is the velocity.

The thermodynamic properties of substances in standard states are temperature "ideal", therefore these properties are defined as temperature-dependent functions and at a constant pressure the heat capacity of these substances is written as a function depending on polynomial coefficients and temperature:

$$\frac{C_{pk}^0}{R} = \sum_{n=1}^N a_{nk} T_k^{(n-1)} \quad (7)$$

Here the sign of the superscript "0" denotes the properties of certain substance in 1 atm. pressure. For ideal gases the heat capacity is independent of temperature, so the value of the heat capacity in the standard state is relevant for them. Other thermodynamic properties of substances at a given temperature are determined in the form of integral functions depending on the specific heat.

The molar enthalpy of substances in standard states is determined by the following formula:

$$H_k^0 = \int_{298}^{T_k} C_{pk}^0 dT + H_k^0(298) \quad (8)$$

$$\frac{H_k^0}{RT_k} = \sum_{n=1}^N \frac{a_{nk} T_k^{(n-1)}}{n} + \frac{a_{N+1,k}}{T_k} \quad (8.1)$$

here, parametre $a_{N+1,k} * R$ denotes the heat of formation at 298K.

The molar entropy in the standard state is written in the form

$$S_k^0 = \int_{298}^{T_k} \frac{C_{pk}^0}{T} dT + S_k^0(298) \quad (9)$$

$$\frac{S_k^0}{R} = a_{1k} \ln T_k + \sum_{n=2}^N \frac{a_{nk} T_k^{(n-1)}}{(n-1)} + a_{N+2,k} \quad (9.1)$$

here, the constant of integration $a_{N+2,k} * R$ is determined depending on the value of the entropy of a given substance at a temperature of 298 K.

The equations, which are given above, are indicated for polynomial of the permissible limits, used in the NASA chemical equations (formula 8). In the Chemkin software complex, these components are divided into two groups for two intervals of the temperature interval (low and high temperature region). In this case, to process the thermodynamic properties [11-14]:

$$\frac{C_{pk}^0}{R} = a_{1k} + a_{2k} T_k + a_{3k} T_k^2 + a_{4k} T_k^3 + a_{5k} T_k^4 \quad (10)$$

$$\frac{H_k^0}{RT_k} = a_{1k} + \frac{a_{2k}}{2} T_k + \frac{a_{3k}}{3} T_k^2 + \frac{a_{4k}}{4} T_k^3 + \frac{a_{5k}}{5} T_k^4 + \frac{a_{6k}}{6} \quad (11)$$

$$\frac{S_k^0}{R} = a_{1k} \ln(T_k) + a_{2k} T_k + \frac{a_{3k}}{2} T_k^2 + \frac{a_{4k}}{3} T_k^3 + \frac{a_{5k}}{4} T_k^4 + a_{7k} \quad (12)$$

Other thermodynamic properties have values C_p^0 , H^0 и S^0 . Specific heat at constant volume $-C_{vk}^0$

$$C_{vk}^0 = C_{pk}^0 - R \quad (13)$$

Interior energy U_k^0

$$U_k^0 = H_k^0 - RT \quad (14)$$

Gibb's free energy in a standard state G_k^0

$$G_k^0 = H_k^0 - T_k S_k^0 \quad (15)$$

Helmholtz free energy in the standard state A_k^0

$$A_k^0 = U_k^0 - T_k S_k^0 \quad (16)$$

For ideal gases in the standard state, the internal heat capacity, enthalpy and specific heat are equal to the actual value of these properties.

Equations for determining the reaction's rate

$$\sum_{k=1}^K v'_{ki} \chi_k \Leftrightarrow \sum_{k=1}^K v''_{ki} \chi_k \quad (i = 1, \dots, I) \quad (17)$$

Stoichiometric coefficients v_{ki} integers and χ_k chemical symbol of k —substance. v'_{ki} — direct reaction's rate, v''_{ki} —reverse reaction's rate. Usually, three or four substances participate in elementary reactions.

To calculate the reaction rates in the "input data" area is set the initial temperature of the mixture and by clicking on the menu item **Run** → **Calculate velocities**, for each reaction, the rates of the forward and reverse reaction are calculated.




Figure 4 - Running the calculation on the GPU

In this program, the basic computational work can be done with the use of a sequential algorithm on the central processor as well as with the use of parallel algorithm in the graphics processor. Before calculating is chosen a method for calculating the change in concentration over time. Then, the time step and the number of cycles are set. The total calculation time is determined as follows:

$$t = \text{Time step} \times \text{Number of cycles}$$

As it can be seen from the input data, to calculate the change of concentration of all elements of this mechanism a sufficiently resource-intensive calculation is required. With a sufficiently large volume of the computation mechanism, using a sequential algorithm requires a lot of computational costs and time. And the use of a parallel algorithm makes it possible to speed up the computation with increasing the volume of the mechanism [2; 4; 11-14].




Figure 5 - Checking the concentration value before starting the calculation

For further calculation, the initial concentration of reagents in the mechanism is given in the "Concentration of substances" area. If the initial concentration for some substances is not specified, the initial concentration for these reagents is taken as zero (figure 5). After the end the calculation results are displayed in the form of a table (figure 6). For clarity, you can see this data in the form of a graph (figure 7). To do this, just choose.





Figure 6 - Results of calculation

menu item **Result** → **Show res. GPU / CPU** → **Show in the graph**. After that, a separate window opens with the graphics area and in the right-hand area of the names of the reagent's mechanism and a label. At the inserting / removing a check mark in these marks a graph concentration changes of the substance over time appears in the graph area [17; 22; 24-27].



орталық процессорда тізбекті түрде есептеледі және параллельді есептеу үшін CUDA графикалық процессорларда параллельді есептеу платформасы қолданылды. Берілген уақыт аралығында белгілі заттың концентрациясының өзгеруін GPU-да деректерді параллельді есептеу үшін реакцияларды бөлек бағыттарға бөледі. Әр бағыт берілген уақыт аралығында белгілі заттың концентрациясы осы реакцияларда бөлінуі әр бағытта есептеледі. Параллельдеу барлық қарапайым реакциялар үшін жасалғандықтан, механизмдегі реакциялар санының артуымен GPU есептеу уақытты әлде-қайда ұтымды етуге алып келеді.

Түйін сөздер: CUDA, жану, модельдеу, параллельді есептеу, өнімділік, есептеу, термодинамика, концентрацияның өзгеруі, механизм реакциялары.

М. С. Сарсембаев¹, Б. А. Урмашев¹, О. Ж. Мамырбаев², М. Турдалыұлы³, Т. С. Сарсембаева¹

¹Казахский национальный университет имени аль-Фараби, Алматы, Казахстан;

²Институт информационных и вычислительных технологий, Алматы, Казахстан;

³Сатпаев Университет, Алматы, Казахстан

ПРИМЕНЕНИЕ ТЕХНОЛОГИИ CUDA ДЛЯ УСКОРЕНИЯ ПРОБЛЕМ ХИМИЧЕСКОЙ КИНЕТИКИ

Аннотация. Основная идея реализации - сократить время вычислений и тем самым реализовать многопользовательский режим для пользователей, разместив его на сервере с доступом через веб-браузер. Для моделирования кинетики химических реагирующих систем использованы методы Рунге – Кутты 4-го и 5-го порядка. С помощью этих методов было написано код программы на языке C# для последовательного вычисления на центральном процессоре и использовано платформа для параллельного вычисления CUDA на графических процессорах. Распараллеливания данных при расчете на GPU производится распределением реакции на отдельные нити, при вычислении изменений концентрации на заданном временном интервале определенного вещества. Каждая нить вычисляет, насколько изменилась концентрация данного вещества в данной реакции на заданном временном интервале. Так как распараллеливание производится по всем элементарным реакциям, с увеличением количеств реакций в механизме вычисление на GPU дает заметный выигрыш по времени.

Ключевые слова: CUDA, горение, моделирование, параллельные вычисления, производительность, вычисления, термодинамика, изменения концентрации, реакции механизма.

Information about authors:

Sarsembayev M., researcher, Al-Farabi Kazakh National University, Almaty, Kazakhstan. magjan@kaznu.kz, <https://orcid.org/0000-0003-2139-2456>;

Urmashiev B., Candidate in Phys. and Math. Sci., Dean of faculty of "Information technologies", Al-Farabi Kazakh National University, Almaty, Kazakhstan. baidaulet.urmashev@kaznu.kz, <https://orcid.org/0000-0002-0085-1708>;

Mamyrbayev O., PhD, Institute of Information and Computational Technologies, Almaty, Kazakhstan. morkenj@mail.ru, <https://orcid.org/0000-0001-8318-3794>;

Turdalyuly M., PhD, Head of Department, Satbayev University, Almaty, Kazakhstan. m.turdalyuly@satbayev.university, <https://orcid.org/0000-0002-1470-3706>;

Sarsembayeva T., department of "Artificial intelligence and big data", Al-Farabi Kazakh National University, Almaty, Kazakhstan. sagdatbek.talshyn1@kaznu.kz, <https://orcid.org/0000-0001-7668-2640>

REFERENCES

- [1] Spatford K., Meredith J., Vetter J., Chen J. H., Grout R., Sankaran R. Accelerating S3D: A GPGPU case study, in: Euro-Par 2009 Parallel Processing Workshops, LNCS 6043, Springer-Verlag, Berlin, Heidelberg, 2010, pp.122–131.
- [2] Burke M. P., Chaos M., Ju Y., Dryer F. L., Klippenstein S. J. Comprehensive H₂/O₂ kinetic model for high-pressure combustion, Int. J. Chem. Kinet. 44 (2011) 444–474.
- [3] Shi Y., Green W. H., Wong H., Oluwole O. O. Accelerating multi-dimensional combustion simulations using hybrid CPU-based implicit/GPU-based explicit ODE integration, Combust. Flame 159 (2012) 2388–2397.
- [4] Chung T.J. Computational Fluid Dynamics. Cambridge University Press, 2002. p. 1012.
- [5] Shi Y., Green W. H., Wong H., Oluwole O. Redesigning combustion modeling algorithms for the graphics processing unit (GPU): Chemical kinetic rate evaluation and ordinary differential equation integration, Combust. Flame 158 (2011) 836–847.
- [6] NVIDIA, CUDA C Programming Guide, 4.0 ed., 2011.
- [7] van der Houwen P. J. The development of Runge–Kutta methods for partial differential equations, Applied Numerical Mathematics 20 (1996) 261–272.
- [8] Sankaran R. GPU-accelerated software library for unsteady flamelet modeling of turbulent combustion with complex chemical kinetics, in: 51st AIAA Aerospace Sciences Meeting, AIAA 2013-0372, 2013.
- [9] Ma F. Performance and emission characteristics of a turbocharged spark-ignition hydrogen-enriched compressed natural gas engine under wide open throttle operating conditions / Ma F., Wang M., Jiang L., Deng J., Chen R., Naeve N., Zhao S. // International journal of Hydrogen Energy. 2010. V. 35. P. 12502–12509.

- [10] Brower M. Ignition delay time experiments and mechanism validation for natural gas/hydrogen blends at elevated pressures / Brower M., Petersen E.L., Metcalfe W., Curran H.J., Füre M., Bourque G., Aluri N., Güthe F. // *J. Eng. Gas. Turb. Power.* – 2013. – ArticleNo. 21504.
- [11] Mehl M., Pitz W. J., Westbrook C. K., Curran H. J. Kinetic modeling of gasoline surrogate components and mixtures under engine conditions, *Proc. Combust. Inst.* 33 (2011) 193–200.
- [12] Belleman R.G., Bédorf J., PortegiesZwart S.F. High performance direct gravitational N-body simulations on graphics processing units II: an implementation in CUDA, 13, *New Astron.*, 2008, pp. 103–112.
- [13] Garland M., S. Le Grand, Nickolls J., Anderson J., Hardwick J., Morton S., Phillips E., Zhang Y., Volkov V. Parallel computing experiences with CUDA, *Micro IEEE* 28 (2008) 13–27.
- [14] Simek V., Dvorak R., Zboril F., Kunovsky J. Towards accelerated computation of atmospheric equations using CUDA, *Proceedings of the UKSim 2009: International Conference on Computer Modelling and Simulation*, 2009, pp. 449–454.
- [15] Dabdub D., Seinfeld J.H. Parallel computation in atmospheric chemical modeling, *Parallel Comput.* 22 (1996) 111–130.
- [16] Senocak I., Thibault J., Caylor M. Rapid-response urban CFD simulations using a GPU computing paradigm on desktop supercomputers, *Eighth Symposium on the Urban Environment*, Phoenix Arizona, 2009, p. J19.2.
- [17] Liu W., Schmidt B., Voss G., Müller-Wittig W. Accelerating molecular dynamics simulations using Graphics Processing Units with CUDA, *Comput. Phys. Commun.* 179 (2008) 634–641.
- [18] Li S., Livshitz B., Lomakin V. Fast evaluation of Helmholtz potential on graphics processing units (GPUs), *J. Comput. Phys.* 229 (2010) 8463–8483.
- [19] Lovas R., Kacsuk P., Lagzi I., Turányi T. Unified development solution for cluster and Grid computing and its application in chemistry, *Lect. Notes Comput. Sci.* 3044 (2004) 226–235.
- [20] Law C. K. *Combustion Physics*, Cambridge University Press, New York, 2006.
- [21] Humphrey J. R., Price D. K., Spagnoli K. E., Paolini A. L., Kelmelis E. J. CULA: Hybrid GPU accelerated linear algebra routines, in: E. J. Kelmelis (Ed.), *Proc. SPIE*, volume 7705, 2010, p. 770502. doi:10.1117/12.850538.
- [22] Kirk D. B., Hwu W. W. *Programming Massively Parallel Processors: A Hands-on Approach*, Morgan Kaufmann, Burlington, MA, 2010.
- [23] J. G. Verwer, W. Hundsdorfer, B. P. Sommeijer, Convergence properties of the Runge–Kutta–Chebyshev method, *Numer. Math.* 57 (1990) 157–178.
- [24] J. G. Verwer, Explicit Runge–Kutta methods for parabolic partial differential equations, *Appl. Numer.* 22(1996) 359–379.
- [25] G. P. Smith, D. M. Golden, M. Frenklach, N. W. Moriarty, B. Eiteneer, M. Goldenberg, C. T. Bowman, R. K. Hanson, S. Song, W. C. Gardiner Jr., V. V. Lissianski, Z. Qin, *GRI-Mech 3.0*, http://www.me.berkeley.edu/gri_mech/, 2010.
- [26] A. Abdule, Fourth order Chebyshev methods with recurrence relation, *SIAM J. Sci. Comput.* 23 (2002) 2041–2054.
- [27] C. P. Stone, R. L. Davis, B. Sekar, Techniques for solving stiff chemical kinetics on GPUs, in: *51st AIAA Aerospace Sciences Meeting*, AIAA 2013-0369, 2013.
- [28] A. C. Zambon, H. K. Chelliah, Explicit reduced reaction models for ignition, flame propagation, and extinction of C₂H₄/CH₄/H₂ and air systems, *Combust. Flame* 150 (2007) 71–91.
- [29] K. E. Niemeyer, C. J. Sung, Recent progress and challenges in exploiting graphics processors in computational fluid dynamics, Manuscript submitted for publication, 2013.

Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

(Правила оформления статьи для публикации в журнале смотреть на сайтах:

www.nauka-nanrk.kz

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы: *М. С. Ахметова, Д. С. Аленов, Р.Ж. Мрзабаева*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 15.04.2021.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
11,6 п.л. Тираж 300. Заказ 2.